

EFFECT OF MUTAGENS ON POLLEN FERTILITY IN M_2 GENERATION OF MUNGBEAN (*Vigna radiata*) (L.) WILCZEK

SHAMIM AHMED AZAD¹

Department of Botany, Govt. Post-Graduate College, Rajouri, Jammu and Kashmir State, India

ABSTRACT

A study was carried out to see the effect of chemical mutagens i.e sodium azide (SA), Hydrazine hydrate (HZ) and Ethyl methane sulphonate (EMS) in the two varieties of Mungbean viz. Asha and K-851 in M_2 generation. The pollen fertility was dose dependent as evident from a proportionate decrease in fertility with the increasing concentrations of mutagens in both the varieties. The highest percentage of reduction was recorded in EMS followed by HZ and SA.

KEYWORDS : Mutagens, K-851, Asha, Pollen Fertility, M_2 Generation

Mungbean is a self-pollinated crop; mutation breeding is an important sources of creating genetic variability. Induced mutation provided a modern and fruitful tool in crop plants for creating genetic variability (Swaminathan, 1969; Gottschalk, 1972; Khan, 1988).

Success of plant breeding including chemo-mutagenesis is directly dependent on the genetic variability in the source population. The major break through in plant breeding was made when man realized that main source of genetic variability was gene mutation and recombination.

Most of the plant attributes of interest to a plant breeder are qualitative Characters which are controlled by polygenic interaction. In such situation the efficiency of selecting the desire mutant is generally lower than for specific characters which are controlled by a single gene. Micromutations produce genetic variability in qualitative characters of the crop plants. Hence, they deserve full attention of plant-breeders. Such mutation should be useful for improving qualitative inherited traits (e.g. grain yield) without disturbing the major part of the genotypic and phenotypic architecture of crop.

In the present study three different chemical mutagens and two varieties of mungbean were taken to see the effect on pollen fertility. The pollen fertility is one of the most important, stable and genetically controlled character which may be considered for hybridization as well as to study the cytological effects to assess the effectiveness and efficiency of mutagens.

MATERIALS AND METHODS

Variety Asha

This variety was released in 1991 for general

cultivation in irrigated areas of Haryana state. The variety is especially suitable for kharif season and semi-erect in growth habit. The seeds are medium, smooth and shining in colour . It matures in 70-75 days. Average yield is 9-11 q /ha.

Variety K-851

The variety K-851 has been developed at Kanpur. It is erect and semi-tall. The seeds are medium bold, smooth and shining green. It matures in 65-70 days. Average yield is 10-12 q/ha.

Mutagens Used

i Ethylmethane sulphonate (EMS)- $CH_3SO_2OC_2H_5$.

It is a monofunctional alkylating agent causes depurination, transition and formation of triesters in the backbone of DNA molecule.

ii Sodium azide (SA) NAN_3 .

It is used as respiratory inhibitor. During duplication of DNA by base transition mechanism, it cause point mutation.

iii Hydrazine hydrate (HZ) $NH_2NH_2.H_2O$.

It is base analogue of nucleic acid and thereby causing gene mutation in the DNA molecule.

M_2 generation

For raising M_2 generation, 50 healthy seeds of both the varieties from each normal looking M_1 plants of all different treatments with their respective control were planted in the plant progeny rows. The different treatments and control comprised of 30 progenies. The spacing was maintained at 30cm plant to plant in a rows and 60cm between the rows in the field. Three replications were maintained in each treatment as well as in control.

¹Corresponding author

Pollen Fertility

The pollen fertility was determined by staining the pollen grains with 1% acetocarmine solution. For this purpose 10 or 15 plants at random were selected from each treatment including their controls for both the varieties and 5 young flower buds from each plant were used for microscopic analysis. Pollen grains which took a deep stain and had a regular outline were considered as fertile, while the shrunken, empty and unstained ones as sterile.

The following formula was used to calculate the percentage of pollen fertility, inhibition or injury or reduction.

Percentage inhibition

or

Percentage injury = $\frac{\text{Control} - \text{treated}}{\text{Control}} \times 100$

or

Percentage reduction

RESULTS

The pollen fertility was dose-dependent as evident from a proportionate decreases in fertility with the

increasing concentrations of mutagens in both the varieties of mungbean in M₂ generation (Table, 1). But it is quite less than the M₁ generation because in M₂ generation the pollen sterility showed a considerable recovery. The highest percentage of reduction was recorded in EMS followed by HZ and SA. The fertility was the lowest 87.50% and 86.62% in at 0.4% EMS in the varieties Asha and K-851 respectively. It ranged from 93.25% to 87.50% in variety Asha and 92.56% to 86.62% in variety K-851. The variety K-851 was found to be more sensitive than Asha.

DISCUSSION

Mutagenic sensitivity is an important aspect of artificial induction of mutations. Species differ with respect to their sensitivity to the action of different physical as well as chemical mutagens. In the present study, the mutagenic sensitivity has been assessed by measuring the effects of mutagenic treatments through pollen sterility in the two varieties viz., Asha and K-851 of mungbean with three different chemical mutagens.

It was observed during mutagenic studies that EMS, SA and HZ bring about a reduction in pollen fertility.

Table 1: Effect of Mutagens on Pollen Fertility in M₂ Generation of Mungbean

Treatment	Var. Asha	Var. K-851		
	Pollen fertility	Percentage of reduction	Pollen fertility	percentage of reduction
Control	97.85	-	96.65	-
0.1 %EMS	93.25	4.70	92.56	3.58
0.2 %EMS	91.05	6.94	90.45	5.78
0.3 %EMS	88.57	9.48	89.15	7.13
0.4 %EMS	87.50	10.57	86.62	9.77
0.1 %SA	94.21	3.71	94.26	1.81
0.02 %SA	93.67	4.27	93.79	2.30
0.03 %SA	92.04	5.93	92.35	3.80
0.04 %SA	90.55	7.46	91.06	5.10
0.01 %HZ	93.32	4.62	92.82	3.31
0.02 %HZ	91.84	6.14	91.00	5.20
0.03 %HZ	91.20	6.79	90.27	5.96
0.04 %HZ	90.06	7.96	89.68	6.58

Such reductions were found to be dose dependent. An immediate consequence of the various aberrations in pollen mother cells due to mutagens could be discerned in the microspores in the form of the pollen sterility. The pollen sterility count is a better way and a dependable parameter to find out the mutagenic effects in M₂ generation.

In the present State of study, varying degree of pollen sterility has been observed in different concentration of the mutagens. The order of the amount of pollen sterility induced by the three mutagens is EMS, HZ and SA. Since SA does not induce chromosomal aberrations, its effect on pollen sterility could be attributed to gene mutations expressed as genetic or Zygotic lethality. High pollen sterility is in EMS and HZ treatments indicating that these two chemical mutagens invite a high degree of meiotic abnormalities.

Dose dependent increase in pollen sterility and a positive and significant correlation between chromosomal abnormality and pollen Sterility was reported earlier and in conformity with the present work (Kallo and Das, 1971 ; Nerker , 1977 ; Grover and Tejpal, 1982 ; Khan and Siddiqui 1988, Khan et al., 1994 ; Khan et. al 1998).

REFERENCES

- Gottschalk W., 1972. Combination of mutated genes as an additional tool in plant breeding. In : Induced mutations and plant improvement. Proc. Symp. IAEA, Vienna.
- Grover I.S. and Tejpal S.K., 1980. Induced mutations in greengram. *Genetica polonica*, **20** (4) : 520-22
- Ignacimuthu S. and Babu C.R., 1988. Radio Sensitivity of the wild and cultivated urd and mungbeans. *Indian Journal of Genet.*, **48** (3) : 331-342.
- Kalloo G. and Das K., 1971. Induced interchange in *Pisum*. *Nucleus*, **14** : 156-160.
- Nerker N.S., 1977. Mutagenic effectiveness and efficiency of gamma rays, ethyl methane sulphonate and nitromethyl urea in *Lathyrus sativus*. *Indian Journal of Genet.*, **37** (1): 37 141.
- Swaminathan M.S., 1969. Role of mutation breeding in a Induced mutations in plants. Proc. Symp. Pullman, 1969. IAEA, Vienna : 719-734.
- Khan A. , 1988. Mutation breeding in mungbean. In : Recent Advances in Genetics and cytogenetics, Premier Pub. House, Hyderabad : 91-102.
- Khan S. and Siddiqui B.A. ,1988. MMS Induced variation in green gram (*Vigna radiata* L. wilczek). In : xvith Int. congress of Genetics, Toronto, Canada : 109.
- Khan S., Siddiqui B.A. and Nadeem M., 1994. Variation in quantitative characters of mungbean after seed treatment with DES. *Adv. Plant Sci.*, **7**(1) : 41 45.
- Khan S., Siddiqui B.A., Rehman M.R. and Azad shamim Ahmed, 1998. Response of Green gram (*Vigna radiata* L.) wilczek to maleic hydrazide. *Journal Indian bot. soc.*, **77**: 95-98.